

Nuclear Medicine DaTscan

If you must cancel or reschedule your appointment, please call us at least **48 hours** prior to your appointment time or you may be held responsible for \$2,468.00 for the radioactive tracer.

TO PREPARE FOR YOUR DATSCAN, YOU WILL NEED TO STOP TAKING THE FOLLOWING MEDICATIONS:

- | | |
|---|-----------------------------|
| • Ephedrine, ketamine, isoflurane | Stop 1 day prior to study |
| • Cocaine, methylphenidate..... | Stop 2 days prior to study |
| • Methamphetamine, mazindol, modafinil..... | Stop 3 days prior to study |
| • Benztropine, fentanyl..... | Stop 5 days prior to study |
| • Amphetamine, dexamphetamine..... | Stop 7 days prior to study |
| • Bupropion (Wellbutrin)..... | Stop 8 days prior to study |
| • Phentermine, phencyclidine..... | Stop 14 days prior to study |

THE DAY OF DATSCAN:

- Bring your script, referral/authorization, insurance card and a form of identification.
- List of your current medications.
- Drink 16oz of water one (1) hour prior to the study.
- Wear comfortable clothing.

If you have any questions or concerns about your DaTscan, please call SJRA scheduling department (888) 909-7572 (Option #1), Monday--Friday from 8:00am to 4:30 pm.