Nuclear Medicine Thyroid Uptake & Scan

Bring any recent thyroid blood work or have your doctor's office fax it to the office where your appointment has been scheduled. Voorhees fax# (856) 751-9676 / Turnersville fax# (856) 227-8537

If you must cancel or reschedule your appointment, please call us at least <u>24 hours</u> prior to your appointment time.

PLEASE DO <u>NOT</u> EAT OR DRINK ANYTHING ONE (1) HOUR PRIOR TO YOUR APPOINTMENT TIME

THE FOLLOWING MEDICATIONS WILL AFFECT YOUR THYROID UPTAKE

	MAJOR GROUPS WITH EXAMPLES	RECOMMENDED TIME TO STOP FOR VALID
<u>UPT</u>		STOT FOR VILLED
•	RADIOGRAPHIC CONTRAST (e.g. CT dye)	6
	WEEKS	
•	THYROID MEDICATIONS	
	 THYROID HORMONE-SYNTHROID (LEVOXYL, LEVOTHYRO 	XINE) 6
	WEEKS	,
	o AMOUR THYROID	4
	WEEKS	
	o CYTOMEL	2
	WEEKS	
•	ANTI-THYROID DRUGS	5 DAYS
	PTU (PROPYLTHIURACIL) TARAZOLE (AUTTHMAZOLE)	
	TAPAZOLE (METHIMAZOLE)PERCHLORATE	
	O PERCHLORATE TOPICAL IODINE COMPOUNDS	4
•	WEEKS	4
	ORAL IODIDES & MEDICATION CONTAINING IODINE	4
•	WEEKS	
	LUGOL'S SOLUTION	
	o POTASSIUM IODIDE	
	o KELP	
	 ANTI-PARASITIC DRUGS-IODOCHLORHYDROQUIN 	
	 AMIODARONE HYDROCHLORIDE 	
•	MULTI-VITAMINS ONLY / FISH OIL SUPPLEMENTS	1
	WEEK	
	o INCLUDES <u>ALL</u> NUTRITION DRINKS AND/OR SHAKES	
•	OTHER MEDICATIONS	1
	WEEK	
	ANTIHISTAMINES COLICIL & COLICIL MEDICATIONS	
	COUGH & COLD MEDICATIONSBUTAZOLIDIN	
	 BUTAZOLIDIN ACTH & CORTICOSTEROIDS 	
	SULFONAMIDES	
•	SEAFOOD (Any Type)	3
_	DAYS	······································

BEFORE STOPPING ANY MEDICATION PLEASE CHECK WITH YOUR PHYSICIAN

If you have any questions or concerns, please contact SJRA scheduling department at (888) 909-7572 option 1, Monday – Friday 8am – 4:30pm